

AD-A125 580

A NOTE ON 'GEOMETRIC TRANSFORMS' OF DIGITAL SETS(U)

MARYLAND UNIV COLLEGE PARK COMPUTER VISION LAB

A ROSENFELD SEP 82 TR-1218 AFOSR-TR-83-0066

AFOSR-77-3271

UNCLASSIFIED


F/G 12/1

NL



END
DATE
FILMED
4 83
DTIC

M-2



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

4

AD A125580

TR-1218
AFOSR-77-3271

September 1982

A NOTE ON "GEOMETRIC
TRANSFORMS" OF DIGITAL SETS

Azriel Rosenfeld

Computer Vision Laboratory
Computer Science Center
University of Maryland
College Park, MD 20742

COMPUTER SCIENCE
TECHNICAL REPORT SERIES

DTIC FILE COPY



Approved for public release;
distribution unlimited.

UNIVERSITY OF MARYLAND
COLLEGE PARK, MARYLAND

20742

DTIC
ELECTE
MAR 14 1983

B

88 03 14 026

UNCLASSIFIED

TR-1218
AFOSR-77-3271

September 1982

A NOTE ON "GEOMETRIC
TRANSFORMS" OF DIGITAL SETS

Azriel Rosenfeld

Computer Vision Laboratory
Computer Science Center
University of Maryland
College Park, MD 20742

DTIC
ELECTE
MAR 14 1983

B

Document defines ABSTRACT

We define a "geometric transform" on the digital plane as a function f that takes pairs (P, S) , where S is a set and P a point of S , into nonnegative integers, and where $f(S, P)$ depends only on the positions of the points of S relative to P . Transforms of this type are useful for segmenting and describing S . Two examples are "distance transforms," for which $f(S, P)$ is the distance from P to S , and "isovist transforms," where $f(S, P)$ is (e.g.) the area of the part of S visible from P . This note characterizes geometric transforms that have certain simple set-theoretic properties, e.g., such that $f(S \cap T, P) = f(S, P) \wedge f(T, P)$ for all S, T, P . It is shown that a geometric transform has this intersection property if and only if it is defined in a special way in terms of a "neighborhood base"; the class of such "neighborhood transforms" is a generalization of the class of distance transforms.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE: This report is the property of the Air Force Office of Scientific Research and is loaned to your organization. It and its contents are not to be distributed outside your organization.
THIS REPORT IS AVAILABLE FROM:
DISTRIBUTION STATEMENTS
MATTHEW J. ...
Chief, Technical Information Division

The support of the U.S. Air Force Office of Scientific Research under Grant AFOSR-77-3271 is gratefully acknowledged, as is the help of Janet Salzman in preparing this paper.

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

1. Introduction

Given a subset S of a digital picture, there are various useful ways of defining functions on S that associate with each point P of S some geometric property of S "relative to P ". An early example [1] is the distance transform, which associates with each $P \in S$ the distance (with respect to some given metric) from P to \bar{S} (the complement of S). This transform is a useful tool for describing or segmenting S ; for example, the well-known "medial axis transformation" of S is just the set of local maxima of its distance transform. A more recent example [2] is the class of "isovist transforms", which associate with each P some property of the part of S "visible" from P , e.g., its area; such transforms can be used, e.g., to find minimal sets of points from which all of S can be seen. (A point Q of S is said to be visible from P if the straight line segment \overline{PQ} lies entirely in S .)

In this note we give a general definition of such "geometric transforms" (for brevity: G-transforms). We also characterize G-transforms that have certain simple properties with respect to set-theoretic operations. In particular, we consider G-transforms having the "intersection property": for any two sets S and T , the transform values for $S \cap T$ are (pointwise) the infs of the values for S and for T . We show that a G-transform has this property iff it can be defined in a special way in terms of a "neighborhood basis"; the class of such transforms includes the class of distance transforms. Interestingly, the analogously defined "union property" implies that the transform must be trivial.

2. G-transforms

Let Σ be a bounded set of lattice points in the plane (e.g., a digital picture), let 2^Σ be the set of subsets of Σ , and let f be a function defined on $2^\Sigma \times \Sigma$. For simplicity, we shall assume that f is integer-valued; that $f(S,P)=0$ whenever $P \notin S$; and that $f(S,P) > 0$ whenever $P \in S$. We call f a G-transform if $f(S,P)$ depends only on the positions of the (other) points of S relative to P . This is a rather general definition; the following are a few examples of G-transforms:

- a) The characteristic function, i.e., $f(S,P)=1$ iff $P \in S$
- b) The distance transform, i.e., $f(S,P)=$ the distance from P to \bar{S}
- c) The "area transform": $f(S,P)=$ the area of the connected component of S that contains P
- d) The isovist transform: $f(S,P)=$ the area of the part of S visible from P

Since a G-transform is defined in terms of positions relative to P , it is evidently shift-invariant -- in other words, shifting S cannot change the G-transform values of its points.* In particular, we have

Proposition 1. $f(\{P\},P)$ has the same value for any P . ||

For simplicity, we assume that this value is 1.

*We assume that when S shifts, it remains inside Σ . Alternatively, we could allow cyclic shifts, and define $f(S,P)$ in terms of the positions of the points of S relative to P "modulo Σ ".

We say that f has the union property if $f(S \cup T, P) = f(S, P) \vee f(T, P)$ for all S, T, P , and the intersection property if $f(S \cap T, P) = f(S, P) \wedge f(T, P)$ for all S, T, P . Evidently the characteristic function has both the union and the intersection property. In fact, it is the only G-transform that has the union property, as we see from Proposition 2. A G-transform f has the union property iff it is the characteristic function.

Proof: By Proposition 1, $f(\{P\}, P) = 1$ for all P . It follows from the union property that $f(\{P, Q\}, P) = f(\{P\}, P) \vee f(\{Q\}, P) = 1$ for all $\{P, Q\}$, i.e., for any two-element subset of Σ . By induction, the same is true for any finite subset of Σ . ||

The G-transforms that have the intersection property are less trivial; we shall characterize them in the next section.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Approved for release	
Dist	Grade
A	

3. N-transforms

Let $n: \{O\} = N_0 \subset N_1 \subset N_2 \subset \dots$ be a nested set of finite subsets of Σ that contain the origin O . For any point P , let N_{pi} be the result of shifting N_i to bring O into the position of P ; thus $n_p: \{P\} = N_{p0} \subset N_{p1} \subset N_{p2} \subset \dots$ is a nested set of sets that contain P . We call n_p a neighborhood basis for P .

Let $1 = n_0 \leq n_1 \leq n_2 \leq \dots$ be any monotonic nondecreasing sequence of positive integers. For any $S \in 2^\Sigma$ and any $P \in S$, there is a largest i , call it $i(S, P)$, such that $N_{pi} \subseteq S$. (Note that $N_{p0} = \{P\} \subseteq S$, and that S is finite.) Let the G-transform f be defined by $f(S, P) = n_{i(S, P)}$. We call such a G-transform an N-transform.

It is easily verified that a distance transform is a N-transform. In fact, let N_i be the "disk" of radius i centered at O , i.e., the set of points whose distances from O are $\leq i$, and let $n_i = i + 1$; then the distance transform $f(S, P)$ is just n_{pi} (1 greater than the radius of the largest disk centered at P and contained in S). Note also that the characteristic function is an N-transform, if we simply take $n_i = 1$ for all i .

Theorem 3. A G-transform f has the intersection property iff it is an N-transform.

Proof: For any S and T we have $i(S \cap T, P) = i(S, P) \wedge i(T, P)$, since the N_p 's are nested. Thus if f is an N-transform we have $f(S \cap T, P) = n_{i(S, P) \wedge i(T, P)} = n_{i(S, P)} \wedge n_{i(T, P)}$ (since the n 's are monotonic) $= f(S, P) \wedge f(T, P)$, so that f has the intersection property.

Conversely, let f be a G-transform and have the intersection property. For any k , if $f(S,P)=f(T,P)=k$, we have $f(S \cap T, P)=k$; thus if there are any sets S such that $f(S,P)=k$, there is a smallest such set, call it S_{Pk} . By shift invariance, $f(S,P)=k$ implies $f(S',P')=k$, where S' is S shifted to make P coincide with P' ; thus $S_{P',k}$ exists iff S_{Pk} does, and they are translates of one another. Let $1=k_0 < k_1 < \dots$ be those k 's for which S_{Pk} exists; then $n_P: \{P\} = N_{P0} \subset N_{P1} \subset \dots$, where $N_{Pi} = S_{Pk_i}$, is a neighborhood basis for P . Moreover, for any S , let $i(S,P)$ be the largest i such that $N_{Pi} \subset S$, and let $f(S,P)=m$. If we had $m=k_j > k_i$, S would have to contain $S_{Pk_j} = N_{Pj}$, contradicting the definition of i . On the other hand, if $m=k_h < k_i$, by the intersection property $k_j = f(N_{Pi}, P) = f(S \cap N_{Pi}, P) = f(S, P) \wedge f(N_{Pi}, P) = k_h$, contradiction. Hence $f(S,P)=k_i$, so that f is an N-transform. \parallel

Thus we see that the intersection property characterizes a class of G-transforms that constitute a natural generalization of the distance transforms.

4. Concluding remarks

The main result of this note has been a "set-theoretic" characterization of the "distance-like" G-transforms. It would be of interest to develop characterizations of other useful classes of G-transforms.

References

1. A. Rosenfeld and J. L. Pfaltz, Sequential operations in digital picture processing, J. ACM 13, 1966, 471-494.
2. L. S. Davis and M. L. Benedikt, Computational models of space: Isovists and isovist fields, Computer Graphics Image Processing 11, 1979, 49-72.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 83-0066	2. GOVT ACCESSION NO. AD-4125 580	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A NOTE ON "GEOMETRIC TRANSFORMS" OF DIGITAL SETS		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) A. Rosenfeld		6. PERFORMING ORG. REPORT NUMBER TR-1218
9. PERFORMING ORGANIZATION NAME AND ADDRESS Computer Vision Laboratory Computer Science Center University of Maryland College Park, MD 20742		8. CONTRACT OR GRANT NUMBER(s) AFOSR-77-3271
11. CONTROLLING OFFICE NAME AND ADDRESS Math. & Info. Sciences, AFOSR/NM Bolling AFB Washington, DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61142F 2344/A2
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE September 1982
		13. NUMBER OF PAGES 7
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Image Processing Pattern recognition Geometric transforms Distance transforms		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We define a "geometric transform" on the digital plane as a function f that takes pairs (P, S), where S is a set and P a point of S, into nonnegative integers, and where $f(S, P)$ depends only on the positions of the points of S relative to P. Transforms of this type are useful for segmenting and describing S. Two examples are "distance transforms," for which $f(S, P)$ is the distance from P to \bar{S}, and "isovist transforms," where $f(S, P)$ is (e.g.) the area of the part of S visible from P. This note characterizes geometric transforms that		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

have certain simple set-theoretic properties, e.g., such that $f(S \cap T, P) = f(S, P) \cap f(T, P)$ for all S, T, P . It is shown that a geometric transform has this intersection property if and only if it is defined in a special way in terms of a "neighborhood base"; the class of such "neighborhood transforms" is a generalization of the class of distance transforms.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)